arXiv:math/0305315v1 [math.AT] 22 May 2003

SELF MAPS OF HP^n VIA THE UNSTABLE ADAMS SPECTRAL SEQUENCE

GUSTAVO GRANJA

Abstract. We use obstruction theory based on the unstable Adams spectral sequence to construct self maps of finite quaternionic projective spaces. As a result, a conjecture of Feder and Gitler regarding the classification of self maps up to homology is proved in two new cases.

1. Introduction

Let $\mathbb{H}\mathrm{P}^n$ denote n-dimensional quaternionic projective space and recall that $H^*(\mathbb{H}\mathrm{P}^n;\mathbb{Z}) = \mathbb{Z}[u]/u^{n+1}$ with $|u| = 4$. Given a self map f of $\mathbb{H}\mathrm{P}^n$, the *degree* of f is the integer $\deg(f)$ such that $f^*(u) = \deg(f)u$.

The homotopy classification of self maps of $\mathbb{H}\mathrm{P}^\infty$ is well known: self maps are classified by their degrees and the allowable degrees are zero and the odd square integers [Ms]. The situation for finite projective spaces is more complicated. It is not true in general that self maps are classified by their degrees [MR], and even the set of possible degrees is unknown. We will write

$$\mathbf{R}_n = \{\deg(f) \in \mathbb{Z}\colon f : \mathbb{H}\mathrm{P}^n \to \mathbb{H}\mathrm{P}^n\}$$

for the set of possible degrees. Alternatively, $\mathbf{R}_n$ can be described as the set of self maps of S^3 which admit an A_n structure (see [St]).

There is a conjectural description of $\mathbf{R}_n$ due to Feder and Gitler. Consider the congruences

$$(1) \qquad C_n : \qquad \prod_{i=0}^{n-1}(k - i^2) \equiv 0 \bmod \begin{cases} (2n)! & \text{if } n \text{ is even,} \\ \frac{(2n)!}{2} & \text{if } n \text{ is odd.} \end{cases}$$

and define the sets of integers

$$\mathbf{FG}_n = \{k \in \mathbb{Z} \quad | \quad k \text{ is a solution of } C_1, \ldots, C_n\}.$$

In [FG], Feder and Gitler computed the restrictions imposed by K-theory and Adams operations on the degrees of self maps of $\mathbb{H}\mathrm{P}^n$: they are precisely that the degree should be in $\mathbf{FG}_n$. Therefore $\mathbf{R}_n \subset \mathbf{FG}_n$.

Conjecture 1.1 (Feder and Gitler). $\mathbf{R}_n = \mathbf{FG}_n$.

This is trivial for $n = 1$. A proof for $n = 2$ is contained in [AC], and for $n = 3$ in [MG2]. Feder and Gitler showed in [FG] that the integers in $\mathbf{FG}_\infty$ are precisely the odd square integers and zero so the conjecture also holds for $n = \infty$. In fact, self maps of $\mathbb{H}\mathrm{P}^\infty$ with these degrees were first constructed by Sullivan in [Su], and

---

Key words and phrases. Self maps, quaternionic projective space, unstable Adams spectral sequence. Mathematics Subject Classification 2000: 55S35,55S36,55S37.

Research supported in part by grants from FCT, FLAD and Fundação Gulbenkian.

it was this which motivated Feder and Gitler's work. McGibbon has also proven that the conjecture holds stably, in a suitable sense (see [MG1, Theorem 3.5]).

The solutions to the congruences (1) are more easily described one prime at a time. It turns out (see Proposition 2.1) that the p-local integers which satisfy $C_1, \ldots, C_n$ consist of the p-adic squares (if $p = 2$ they must also be units or zero) together with all multiples of $p^{e(p,n)}$ for a certain function e described in (2). The former can all be realized as degrees of self maps of the p-localization $\mathbb{HP}^\infty_{(p)}$ by work of Rector (see Proposition 2.2), so to prove the conjecture it is enough to build self maps of $\mathbb{HP}^n_{(p)}$ with degree an arbitrary multiple of $p^{e(p,n)}$.

The obstruction to extending a self map of $\mathbb{HP}^n$ to $\mathbb{HP}^{n+1}$ is an element in $\pi_{4n+2}S^3$ (see (5)). We will use some of the information available on $\pi_* S^3$ to analyse this obstruction and produce self maps of $\mathbb{HP}^n_{(p)}$ with degree any multiple of $p^{f(p,n)}$ where f is a different function described in (11) (see Theorem 4.2). In general, $e(p,n) < f(p,n)$ but for $n \leq 5$ the two functions agree, and this implies our main result.

Theorem 1.2. *The Feder-Gitler conjecture holds for $n \leq 5$.*

This is stronger evidence for the conjecture as the obstruction group $\pi_{4n-2}S^3$ is detected by K-theory and Adams operations (i.e. by the e-invariant) only for $n \leq 3$. For $n \geq 3$ it is easy to produce self maps of $\mathbb{HP}^n$ with degree in $\mathbf{FG}_{n+1}$ which do not extend one stage. Theorem 1.2 is proved by inductively building maps of high Adams filtration thus ensuring their obstructions to extension are detected by the e-invariant.

Organization of the paper. In section 2 we describe the p-local solutions to the congruences (1) and quote results of Rector which reduce the Feder-Gitler conjecture to producing maps with degree any multiple of $p^{e(p,n)}$. In section 3, we gather the necessary information on the unstable Adams spectral sequence for S^3. In section 4 we show that the degree of a self map of $\mathbb{HP}^n$ is in $\mathbf{FG}_{n+1}$ if and only if the Adams e-invariant of the obstruction to extension vanishes. The results quoted in section 3 then allow us to prove Theorem 1.2.

Acknowledgments. The author thanks Mike Hopkins for his guidance and useful suggestions, and also Maxence Cuvilliez for a useful discussion.

2. THE LOCAL VERSION OF THE CONJECTURE

We will write $|k|_p = \sup\{l \colon p^l | k\}$ for the p-adic valuation of k and $[x]$ for the least integer $\leq x$. In this section we reduce the Feder-Gitler conjecture to constructing self maps of the localizations $\mathbb{HP}^n_{(p)}$ of degree k for all k with $|k|_p \geq e(p,n)$ for a certain function $e(p,n)$.

First note that, by celullar approximation, the inclusion of $\mathbb{HP}^n$ in $\mathbb{HP}^\infty$ induces a bijection $[\mathbb{HP}^n, \mathbb{HP}^n] \to [\mathbb{HP}^n, \mathbb{HP}^\infty]$. As $\mathbb{HP}^\infty_{(0)} = K(\mathbb{Q}, 4)$, it follows easily from Sullivan's arithmetic square that in order to produce a self map of $\mathbb{HP}^n$ of degree k, it suffices to do so after localizing at each prime p (or, in fact, at each prime $p \leq 2n-1$).

We will write $\mathbf{R}_{n,p} = \{\deg(f) \in \mathbb{Z}_{(p)} | f : \mathbb{HP}^n_{(p)} \to \mathbb{HP}^n_{(p)}\}$ and $\mathbf{FG}_{n,p}$ for the set of solutions in $\mathbb{Z}_{(p)}$ of the congruences (1). Let $\mathbf{D}_p$ denote the set defined by

$$\mathbf{D}_p = \begin{cases} \{k \in \mathbb{Z}_{(2)} \mid k = 0 \text{ or } k \equiv 1 \bmod 8\} & \text{if } p = 2, \\ \{k \in \mathbb{Z}_{(p)} \mid k = u^2 \text{ for some } u \in \hat{\mathbb{Z}}_p\} & \text{if } p > 2. \end{cases}$$

It will soon emerge that $\mathbf{D}_p = \mathbf{R}_{\infty,p} = \mathbf{FG}_{\infty,p}$. Note that $\mathbf{D}_2$ consists of the 2-adic unit squares in $\mathbb{Z}_{(2)}$ together with 0. We define the function

$$(2) \quad e(p,n) = \begin{cases} 0 & \text{if } n = 1, \\ \#\{k \leq n \mid k = p^j \text{ or } k = (\frac{p+1}{2})p^{j-1} \text{ for some } j \geq 1\} & \text{if } p > 2, \\ 1 + 2[\log_2(n)] & \text{otherwise.} \end{cases}$$

Proposition 2.1. *Let p be a prime and $n < \infty$. Then*

$$\mathbf{FG}_{n,p} = \mathbf{D}_p \cup p^{e(p,n)}\mathbb{Z}_{(p)}.$$

Proof. The sets $\mathbf{D}_p$ are the closure of $\mathbf{R}_\infty = \{0, 1, 9, 25, \ldots\}$ in $\mathbb{Z}_{(p)}$ with the p-adic topology. Since the solution set of each congruence is closed we conclude that $\mathbf{D}_p \subset \mathbf{FG}_{n,p}$ for every n, p. We will conclude the proof in the case $p = 2$ (the case when p is odd is similar).

If $k \in \mathbb{Z}_{(2)}$ is a solution of C_2 then either $k \equiv 1 \bmod 8$, or $k = 2^3 l$ for some $l \in \mathbb{Z}_{(2)}$ so it suffices to consider solutions k with $|k|_2 \geq 3$. For these, since $|(4m+2)!/2|_2 = |(4m)!|_2$ it suffices to consider the congruences C_{2m}. So let $|k|_2 \geq 3$. Factoring out units in $\mathbb{Z}_{(2)}$ we see that k is a solution of C_{2m} if and only if

$$\prod_{i=0}^{m-1} (k - (2i)^2) \equiv 0 \bmod 2^{|(4m)!|_2}.$$

Writing $k = 4l$ and noting that $|(4m)!|_2 = 2m + |(2m)!|_2$, this becomes

$$\prod_{i=0}^{m-1} (l - i^2) \equiv 0 \bmod 2^{|(2m)!|_2}$$

so if $k = 4l \in \mathbf{FG}_{2m,2}$ then $l \in \mathbf{FG}_{m,2}$. Since $l \notin \mathbb{Z}^\times_{(2)}$, assuming the result is true for all $n < 2^d$ and letting $2^d \leq 2m < 2^{d+1}$, it follows by induction that $\mathbf{R}_{2m,2} - \mathbf{D}_2 \subset 2^{2d+1}\mathbb{Z}_{(2)}$.

It remains to check that for $2^d \leq 2m < 2^{d+1}$ we have $k \in \mathbf{FG}_{2m,2}$ as long as $|k|_2 \geq 2d + 1$. That is, for $j \leq m$ and $|k|_2 \geq 2d + 1$,

$$\begin{aligned} \left| \prod_{i=0}^{2j-1} (k - i^2) \right|_2 &\geq |(4j)!|_2 \\ |k|_2 + \sum_{i=1}^{2j-1} 2|i|_2 &\geq 2j + j + \ldots + [j/2^{d-1}] \\ |k|_2 + 2(j + \ldots + [j/2^{d-1}]) - 2|2j|_2 &\geq 2j + j + \ldots + [j/2^{d-1}] \\ |k|_2 + j + \ldots + [j/2^{d-1}] &\geq 2j + 2|2j|_2. \end{aligned}$$

The last inequality is easy to check if we write $2j = 2^r v$ with $v \in \mathbb{Z}^\times_{(2)}$ and notice that $r \leq d$ and $2^{d-r} \leq v < 2^{d-r+1}$. □

As $e(p,n)$ tends to ∞ with n, this means that $\mathbf{FG}_{\infty,p} = \mathbf{D}_p$. The following result[1] of Rector [Re] implies that in fact we have $\mathbf{R}_{\infty,p} = \mathbf{FG}_{\infty,p} = \mathbf{D}_p$ and therefore, to prove the conjecture it suffices to construct self maps with degree in $p^{e(p,n)}\mathbb{Z}_{(p)}$.

Proposition 2.2 (Rector). *If $k \in \mathbf{D}_p$ then there is a self map of $\mathbb{H}\mathrm{P}^\infty_{(p)}$ of degree k.*

Remark 2.3. *In view of the above, the Feder-Gitler conjecture implies the following: If a self map of $S^3_{(2)}$ admits an A_{2^k}-structure then it admits an A_n-structure for all $n < 2^{k+1}$ (not necessarily extending the A_{2^k}-structure). Similarly for p odd. It would be interesting to know whether this holds more generally.*

A related question is whether a 2-local H-space admitting an A_{2^k}-structure necessarily admits an A_n structure for $n < 2^{k+1}$. In this regard, note that a grouplike multiplication on a rational space always admits a loop structure [Sch, Corollary 2].

3. The unstable Adams spectral sequence for $\mathbb{H}\mathrm{P}^\infty$

Let $\mathcal{U}$ denote the category of unstable modules over the mod p Steenrod algebra. If $M \in \mathcal{U}$, $U(M)$ denotes the free unstable algebra generated by M and for $P \in \mathcal{U}$ a free module, we write $K(P)$ for the product of mod p Eilenberg-Maclane spaces such that $H^*(K(P);\mathbb{F}_p) = U(P)$. Recall from [HM2] that if X is a simply connected space such that $H^*(X;\mathbb{F}_p) = U(M)$ for some $M \in \mathcal{U}$ of finite type, then given a free resolution

$$0 \longleftarrow M \longleftarrow P_0 \longleftarrow P_1 \longleftarrow P_2 \longleftarrow \dots$$

of M with P_s of finite type and 0 in degree $\leq s+1$, one can construct a tower of principal fibrations over X, called an *Adams resolution for X*

(3)

$$\begin{tikzcd}
\vdots \arrow[d, "p_2"] & \\
X\{2\} \arrow[r, "j_2"] \arrow[d, "p_1"] & K(\Omega^2 P_2) \\
X\{1\} \arrow[r, "j_1"] \arrow[d, "p_0"] & K(\Omega P_1) \\
X\{0\} = X \arrow[r, "j_0"] & K(P_0)
\end{tikzcd}$$

where Ω denotes the algebraic loop functor (the left adjoint to suspension). The corresponding homotopy spectral sequence is the Massey-Peterson version of the unstable Adams spectral sequence. For a simply connected space of finite type this spectral sequence converges strongly and

$$E_2^{s,t}(X) = \mathrm{Ext}^{s,t}_{\mathcal{U}}(M,\mathbb{F}_p) \Rightarrow \pi_{t-s}X \otimes \widehat{\mathbb{Z}}_p.$$

The *Adams filtration* of a map $f : W \to X$ is the largest s such that f factors through $X\{s\}$.

---

[1] See also [JMO] for a more up to date account of self maps of p-completed classifying spaces.

Notice that $H^*(S^3;\mathbb{F}_p) = U(\Sigma^3\mathbb{F}_p)$ and $H^*(\mathbb{HP}^\infty;\mathbb{F}_p) = U(M)$ where M is such that $\Omega M = \Sigma^3\mathbb{F}_p$. One easily checks that the first derived functor $\Omega_1 M = 0$ hence looping an Adams resolution for $\mathbb{HP}^\infty$ yields an Adams resolution for S^3. This in turn implies that the adjunction isomorphim

$$\pi_k S^3 \simeq \pi_k \Omega\mathbb{HP}^\infty \simeq \pi_{k+1}\mathbb{HP}^\infty \tag{4}$$

lifts to an isomorphism of spectral sequences

$$E_r^{s,t}(S^3) = E_r^{s,t+1}(\mathbb{HP}^\infty).$$

We will now collect some well known information about this spectral sequence. Figures 1 and 2 describe the portion of the E_2 term of the spectral sequence for $\mathbb{HP}^\infty$ along the vanishing line (i.e. the v_1-periodic part). As usual, vertical lines represent multiplication by p and the slanted lines composition with η if the prime is 2 and with α_1 if the prime is odd. As usual, for p an odd prime, we set $q = 2(p-1)$.



Figure 1. $E_2^{(s,t)}(\mathbb{HP}^\infty)$ for $p = 2$.



Figure 2. $E_2^{(s,t)}(\mathbb{HP}^\infty)$ for p odd.

Theorem 3.1 (Adams, Mahowald, Miller, Harper-Miller)**.** *Consider Figures 1,2.*

(a) *Above the classes shown and the dotted lines in the columns where no classes appear, the E_2 term vanishes.*

(b) *The classes in dimensions $qk+3$ for p odd and the classes in dimension $8k+7$ with filtration $\geq 4k+2$ for $p = 2$ correspond to elements of $\pi_* S^3$ which are stable and detected by the (stable, real) e-invariant.* [2]

---

[2] See (8) below for the definition of the e-invariant.

Proof. Recall from [Ma1] and [HM1] that for each prime there is a bigraded complex $(\Lambda(3), d)$ such that

$$E_2^{s,t}(S^3) = H^{s,t-3}\Lambda(3)$$

There is a short exact sequence of complexes

$$0 \longrightarrow \Lambda(1) \longrightarrow \Lambda(3) \longrightarrow W(1) \longrightarrow 0$$

which induces a split short exact sequence on homology. $\Lambda(1)$ is a complex with 0 differential which corresponds to the $\mathbb{Z}$-tower in $t - s = 3$.

Let $M \otimes \Lambda$ denote the E_1-term of the stable Adams spectral sequence for the mod p Moore spectrum given by the Λ-algebra. The main results of [Ma1] and [HM1] state that there is a map of complexes $W(1) \longrightarrow M \otimes \Lambda$ inducing an isomorphism on homology for $t - s < 5s - 16$ for $p = 2$ and $t - s < (p+1)qs - (p+2)q$ for p odd. This reduces (a) to a statement about the stable E_2-term of the Moore spectrum except in low dimensions where it is easily checked directly by computing the homology of the complex $\Lambda(3)$.

The E_2-term of the Moore spectrum is computed in the required range in [Mi] for p odd and in [Ma3] for $p = 2$. It agrees with the description given in Figures 1 and 2. This proves (a).

Let M^n denote the mod p Moore space with top cell in dimension n and $i : S^{n-1} \longrightarrow M^n$ denote the inclusion of the bottom cell. Let $A : M^{n+r} \longrightarrow M^n$ with $r = 8$ if $p = 2$ and $r = q$ if $p > 2$ be a map inducing an isomorphism in K-theory (see [Ad]).

Suppose first that p is odd. Let $\tilde{\alpha}$ denote an extension of a generator $\alpha \in \pi_{2p}S^3$ to M^{2p+1}. Then by [Ad, Proposition 12.7], $\alpha_k = \tilde{\alpha} \circ A^{k-1} \circ i \in \pi_{qk+2}S^3$ survives to a stable class which is detected by the e-invariant. Since α_k has Adams filtration at least k it must be represented in $E_2(S^3)$ by the class in bidegree $(t-s, s) = (qk+2, k)$.

Next let $p = 2$. Let $\tilde{\mu}$ denote an extension to M^{13} of the generator μ of $\pi_{12}S^3$. Then $\mu_k = \tilde{\mu} \circ A^{k-1} \circ i \in \pi_{8k+4}S^3$ has Adams filtration at least $4k+1$ (the Adams filtration of the Adams map is 4 since we are in the stable range). It follows that the Toda bracket $< \mu_k, 2, \eta > \in \pi_{8k+7}S^3$ has Adams filtration at least $4k+2$. [Ad, Proposition 12.18] shows that this bracket survives to a stable class of order 4 detected by the e-invariant so the bracket is represented on $E_2(S^3)$ by the class in bidegree $(t-s, s) = (8k+6, 4k+2)$. This completes the proof of (b). □

4. Construction of maps

Let $\nu : S^{4n+3} \to \mathbb{H}\mathrm{P}^n$ denote the Hopf map. The cofiber sequence

$$S^{4n+3} \xrightarrow{\nu} \mathbb{H}\mathrm{P}^n \to \mathbb{H}\mathrm{P}^{n+1}$$

shows that the obstruction to extending a map $f : \mathbb{H}\mathrm{P}^n \to \mathbb{H}\mathrm{P}^\infty$ to $\mathbb{H}\mathrm{P}^{n+1}$ is

$$o(f) := [f \circ \nu] \in \pi_{4n+3}\mathbb{H}\mathrm{P}^\infty. \tag{5}$$

Alternatively, one can regard the obstruction as an element in $\pi_{4n+3}\mathbb{H}\mathrm{P}^{n+1}$, in which case, it is given by the formula

$$o(f) := [f \circ \nu] - \deg(f)^{n+1}[\nu] \in \pi_{4n+3}\mathbb{H}\mathrm{P}^{n+1} \tag{6}$$

as a simple cohomology calculation shows.

Note that, as $S^3 \to \Omega\mathbb{H}\mathrm{P}^\infty$ factors through ΩS^4, the isomorphism (4) shows that any element of $\pi_*\mathbb{H}\mathrm{P}^\infty$ (in particular the obstruction to extension) factors uniquely through the bottom cell of $\mathbb{H}\mathrm{P}^\infty$. We will still write $o(f)$ for the corresponding element in $\pi_{4n+3}S^4$.

We will write $KO(X), K(X), KSp(X)$ for the reduced K-groups of X. Using the Atiyah-Hirzebruch spectral sequence, one easily checks that $K^0(\mathbb{H}\mathrm{P}^n) = \mathbb{Z}[x]/x^{n+1}$, and $KO^0(\mathbb{H}\mathrm{P}^n) = \mathbb{Z}[y,z]/ < y^{n+1}, z - 4y^2 >$ where x and y are the classes represented by the reduced tautological bundle over $\mathbb{H}\mathrm{P}^n$. The complexification map $c : K(\mathbb{H}\mathrm{P}^n) \to KO(\mathbb{H}\mathrm{P}^n)$ sends x to $2y$. The forgetful map $r : K(\mathbb{H}\mathrm{P}^n) \to KO(\mathbb{H}\mathrm{P}^n)$ sends yz^k to x^{2k+1} and z^k to $2x^{2k}$. Moreover $KSp(\mathbb{H}\mathrm{P}^n) = \overline{KO}^4(\mathbb{H}\mathrm{P}^n)$ is a free abelian group of rank n and the image of the forgetful map $KSp(\mathbb{H}\mathrm{P}^n) \to K(\mathbb{H}\mathrm{P}^n)$ consists of those integer combinations of the x^i where the coefficient of x^i is even when i is even.

In [FG], Feder and Gitler show that the Adams operations on $K(\mathbb{H}\mathrm{P}^n)$ are given by the formula

$$\psi^l(x) = l^2 x + \ldots + \frac{2l^2(l^2-1)\ldots(l^2-(n-1)^2)}{(2n)!}x^n. \tag{7}$$

and this in turn determines the action of the Adams operations on $KO(\mathbb{H}\mathrm{P}^n)$ (recall that both c and r commute with Adams operations).

Let $\mathcal{A}$ be Adams' category of abelian groups with Adams operations [Ad, Section 6]. Recall from [Ad] that the (real) e-invariant is a group homomorphism

$$Z \xrightarrow{e} \mathrm{Ext}_{\mathcal{A}}(KO(X), KO(S^{j+1})) \tag{8}$$

where $Z = \{\alpha \in \pi_j(X) | KO(\alpha) = 0\}$. $e(\alpha)$ is the Yoneda class of the extension

$$0 \to KO(S^{j+1}) \to KO(X \cup_\alpha e^{j+1}) \to KO(X) \to 0.$$

If X is a sphere there is a natural identification of the target of e with a subgroup of $\mathbb{Q}/\mathbb{Z}$. The stable e-invariant of $\beta \in \pi_{k+l}S^k$ is defined by representing β as a class $\beta' \in \pi_{8m+l}S^{8m}$ for some m and then setting $e^s(\beta) = e(\beta') \in \mathbb{Q}/\mathbb{Z}$. This is independent of the choice of m.

Proposition 4.1. *Let f be a self map of $\mathbb{H}\mathrm{P}^n$ and let $o(f) \in \pi_{4n+3}S^4$ denote its obstruction to extension. Then $\deg(f) \in \mathbf{R}_{n+1}$ if and only if the stable e-invariant of $o(f)$ vanishes.*

Proof. Let $k = \deg(f)$. Feder and Gitler show in [FG] that $k \in \mathbf{R}_{n+1}$ if and only if there is a ring endomorphism ϕ of $K^0(\mathbb{H}\mathrm{P}^{n+1})$ commuting with Adams operations, such that $\phi(x)$ is in the image of the forgetful map $KSp(\mathbb{H}\mathrm{P}^n) \to K(\mathbb{H}\mathrm{P}^n)$.

In turn, this is equivalent to the existence of a map $\psi : KO(\mathbb{H}\mathrm{P}^{n+1}) \to KO(\mathbb{H}\mathrm{P}^{n+1})$ in $\mathcal{A}$ such that the following diagram in $\mathcal{A}$ commutes

$$\begin{tikzcd}
0 \arrow[r] & KO(S^{4n+4}) \arrow[r] \arrow[d, "k^{n+1}"] & KO(\mathbb{H}\mathrm{P}^{n+1}) \arrow[r] \arrow[d, dashed, "\psi"] & KO(\mathbb{H}\mathrm{P}^n) \arrow[r] \arrow[d, "f^*"] & 0 \\
0 \arrow[r] & KO(S^{4n+4}) \arrow[r] & KO(\mathbb{H}\mathrm{P}^{n+1}) \arrow[r] & KO(\mathbb{H}\mathrm{P}^n) \arrow[r] & 0
\end{tikzcd} \tag{9}$$

and $\psi(y)$ is in the image of the forgetful map $KSp \to KO$.

In fact, a ring endomorphim of $K^0(\mathbb{H}\mathrm{P}^{n+1})$ commuting with Adams operations determines a unique ring endomorphism ψ of $KO^0(\mathbb{H}\mathrm{P}^{n+1})$ commuting with Adams operations and this will necessarily make (9) commute. Conversely, such a ψ is determined by its value $\psi(y)$ since by (7), $\psi^l(y)$ generate $KO^0(\mathbb{H}\mathrm{P}^{n+1})$ over $\mathbb{Q}$. Commutativity of (9) then implies that $\psi(y)$ is determined by $\deg(f)$ and it then follows that $\psi(y)$ must correspond to the unique (see [FG]) endomorphism ϕ of $K^0(\mathbb{H}\mathrm{P}^{n+1})\otimes\mathbb{Q}$ commuting with Adams operations such that $\phi(x) = \deg(f)x + \dots$.

Consider the extension E in $\mathcal{A}$ which is (9) as an extension of groups but where the Adams operations are replaced with $\tilde{\psi}^l = l^2\psi^l$. Because $KO(\mathbb{H}\mathrm{P}^n)$ is torsion free, the existence of the map of extensions (9) is equivalent to the existence of the corresponding self map of E. Cupping with the generator of $KO(S^4)$ yields a map in $\mathrm{Ext}_{\mathcal{A}}$

$$E \longrightarrow (0 \to KO(S^{4n+4}) \to KO(\Sigma^4\mathbb{H}\mathrm{P}^{n+1}) \to KO(\Sigma^4\mathbb{H}\mathrm{P}^n) \to 0) \tag{10}$$

which is termwise an injection of abelian groups. The condition that $\phi(x)$ lie in the image of the forgetful map from KSp is easily seen to be equivalent to the condition that the self map of E extends over the monomorphism (10).

We conclude that $k \in \mathbf{R}_{n+1}$ iff the following self map exists in $\mathrm{Ext}_{\mathcal{A}}$

$$\begin{tikzcd}
0 \arrow[r] & KO(S^{4n+8}) \arrow[r] \arrow[d, "k^{n+1}"] & KO(\Sigma^4\mathbb{H}\mathrm{P}^{n+1}) \arrow[r] \arrow[d, dashed] & KO(\Sigma^4\mathbb{H}\mathrm{P}^n) \arrow[r] \arrow[d, "\Sigma^4 f^*"] & 0 \\
0 \arrow[r] & KO(S^{4n+8}) \arrow[r] & KO(\Sigma^4\mathbb{H}\mathrm{P}^{n+1}) \arrow[r] & KO(\Sigma^4\mathbb{H}\mathrm{P}^n) \arrow[r] & 0
\end{tikzcd}$$

or equivalently, writing $\Sigma^4\nu : S^{4n+3} \to \Sigma^4\mathbb{H}\mathrm{P}^n$ for the attaching map of the top cell of $\mathbb{H}\mathrm{P}^{n+1}$, iff the following equality holds in $\mathrm{Ext}_{\mathcal{A}}$:

$$e(\Sigma^4\nu) \circ \Sigma^4 f^* = k^{n+1} \circ e(\Sigma^4\nu)$$

where $\circ$ denotes the Yoneda product. Since the e-invariant sends compositions of maps to Yoneda products this is the same as $e(\Sigma^4(f \circ \nu)) = e(k^{n+1}\Sigma^4\nu)$ and, since e is a group homomorphism, by (6) this is equivalent to $e(\Sigma^4 o(f)) = 0$ in $\mathrm{Ext}_{\mathcal{A}}(KO(\Sigma^4 HP^n), KO(S^{4n+8}))$.

Since $o(f)$ factors through the inclusion of the bottom cell of $\mathbb{H}\mathrm{P}^n$, letting $X = \Sigma^4\mathbb{H}\mathrm{P}^n \cup_{o(f)} e^{4n+8}$, we have a map of extensions

$$\begin{tikzcd}
0 \arrow[r] & KO(S^{4n+8}) \arrow[r] \arrow[d, "1"] & KO(X) \arrow[r] \arrow[d] & KO(\Sigma^4\mathbb{H}\mathrm{P}^n) \arrow[r] \arrow[d, "i^*"] & 0 \\
0 \arrow[r] & KO(S^{4n+8}) \arrow[r] & KO(S^8 \cup_{\Sigma^4 o(f)} e^{4n+8}) \arrow[r] & KO(S^8) \arrow[r] & 0
\end{tikzcd}$$

where i denotes the inclusion of the bottom cell, so we have the equality

$$e^s(o(f)) \circ i^* = e(\Sigma^4 o(f))$$

The long exact sequence in $\mathrm{Ext}_{\mathcal{A}}$ induced by

$$0 \longrightarrow KO(\Sigma^4(\mathbb{H}\mathrm{P}^n/\mathbb{H}\mathrm{P}^1)) \longrightarrow KO(\Sigma^4\mathbb{H}\mathrm{P}^n) \xrightarrow{i^*} KO(S^8) \longrightarrow 0$$

together with the easily checked fact that $\mathrm{Hom}_{\mathcal{A}}(KO(\Sigma^4(\mathbb{H}\mathrm{P}^n/\mathbb{H}\mathrm{P}^1)), KO(S^{4n+8})) = 0$ show that composition with i^* is injective which concludes the proof. $\square$

We are now ready to complete the proof of our main result. Consider the function

$$f(p,n) = \begin{cases} e(p,n) & \text{for } n \leq 5 \\ 4\left[\frac{n}{2}\right] - 3 & \text{for } p = 2, n > 5 \\ -\left[1 - \frac{2n-2}{p-1}\right] & \text{for } p \geq 3, n > 5. \end{cases} \tag{11}$$

Theorem 1.2 is now an immediate consequence of the following result.

Theorem 4.2. $p^{f(p,n)}\mathbb{Z}_{(p)} \subset \mathbf{R}_{n,p}$.

Proof. Pick an Adams resolution for $\mathbb{HP}^\infty_{(p)}$ constructed from a minimal resolution so that all the d_1-differentials vanish. Then the Adams covers $X\{s\}$ are 3-connected, $\pi_4(X\{s\}) = \mathbb{Z}_{(p)}$ and the projection maps $X\{s\} \to \mathbb{HP}^\infty$ induce multiplication by p^s on π_4. Thus to prove the theorem, it is enough to provide a map $\mathbb{HP}^n \to X\{f(p,n)\}$ of degree k on the bottom cell for an arbitrary $k \in \mathbb{Z}_{(p)}$. This will be obtained by extending a map $\mathbb{HP}^1 \to X\{f(p,n)\}$ of degree k.

Consider first the case when $p = 2$. If n is even, then for $k < n-1$ the obstruction to extension of the map $\mathbb{HP}^k \to X\{f(2,n)\}$ to $\mathbb{HP}^{k+1}$ lies in the group $\pi_{4k+3}X\{f(2,n)\}$. By the vanishing line of Theorem 3.1(a) this group is 0 so the map extends. Thus we have a map $\mathbb{HP}^{n-1} \to X\{f(2,n)\}$. Since the Hopf map $S^{4n-1} \to \mathbb{HP}^{n-1}$ has Adams filtration 1, the composite $S^{4n-1} \to X\{f(2,n)\}$ factors through $X\{f(2,n)+1\}$. Let $\gamma : S^{4n-1} \to X\{f(2,n)+1\}$ denote this obstruction class. By Theorem 3.1(b), its image in $\pi_{4n-1}\mathbb{HP}^\infty_{(2)}$ is detected by the e-invariant (it is in the summand of order 4 corresponding to one of the top lightning flashes). Since the map $\pi_{4n-1}X\{f(2,n)+1\} \to \mathbb{HP}^\infty_{(2)}$ is injective (because there are no nonzero differentials $E_r^{4n,k-r+1} \to E_r^{4n-1,k}$ for $k > f(2,n)$), by Proposition 4.1 we must have $\gamma = 0$. Therefore our map extends to give a map $\mathbb{HP}^n \to X\{f(p,n)\}$ of degree k. Moreover, this map extends one more stage because the obstruction to extension lies above the vanishing line of Theorem 3.1(a). This completes the proof for $p = 2$.

For p odd, the reasoning is identical, except that for $p = 3$ and $n = 5$ we have to use the fact that $\pi_{19}S^3_{(3)}$ is cyclic (see [To]) and hence detected by the e-invariant. □

Remark 4.3. *For p odd, Theorem 4.2 can be improved in a small range of values of n using the following observations:*

(a) For $n < \frac{(2p+1)(p-1)}{2}$ all classes in $\pi_{4n-2}(S^3_{(p)})$ are detected by the e-invariant, therefore the conjecture holds at odd primes for these values of n. This follows from the results of Mahowald and Harper-Miller (cf. proof of Theorem 3.1) together with calculations with the Λ-algebra in low degrees.

(b) Let $g(p,n) = \sup\{k : p^k\mathbb{Z}_{(p)} \subset \mathbf{R}_{n,p}\}$. It is not hard to show that if f, g are self maps of $\mathbb{HP}^n$, the obstruction to extension of the composite is $o(fg) = \deg(g)^n o(f) + \deg(f)o(g)$. Since (by Proposition 2.2) there is a self map of $\mathbb{HP}^n_{(p)}$ of degree p^2, Selick's exponent theorem [Se] *implies that $g(p,n+1) \leq 2 + g(p,n)$.*

References

[Ad] J. F. Adams, *On the groups J(X).IV*, Topology 5 (1966) 21-71.

[AC] M. Arkowitz and C. R. Curjel, *On maps of H-spaces*, Topology 6 (1967) 137-148.

[CM] E. Curtis and M. Mahowald, *The unstable Adams spectral sequence for S^3*, Contemp. Math. 96, AMS (1989) 125-162.

[FG] S. Feder and S. Gitler, *Mappings of quaternionic projective spaces*, Bol. Soc. Mat. Mex. 34 (1975) 12-18.

[Gr] B. Gray, *On the sphere of origin of infinite families in the homotopy groups of spheres*, Topology 8 (1969) 219-232.

[HM1] J. Harper and H. R. Miller, *On the double suspension homomorphism at odd primes*, Trans. Amer. Math. Soc. 273 (1982) 319-331.

[HM2] J. Harper and H. R. Miller, *Looping Massey-Peterson towers* in: S.M. Salamon, B. Steer, W.A. Sutherland, eds. *Advances in Homotopy Theory*(London Math. Soc. Lec. Notes 139, Cambridge University Press, Cambridge, 1989) 69-86.

[JMO] S. Jackowski, J. McClure, and B. Oliver, *Homotopy theory of classifying spaces of compact Lie groups* in: *Algebraic topology and its applications*, 81–123, Math. Sci. Res. Inst. Publ., 27, Springer, New York, 1994.

[Ma1] M. Mahowald, *On the double suspension homomorphism*, Trans. Amer. Math. Soc. 214 (1975) 169-178.

[Ma2] M. Mahowald, *The image of J in the EHP sequence*, Annals of Math. 116 (1982) 65-112.

[Ma3] M. Mahowald, *The order of the image of the J homomorphism*, Bull. Amer. Math. Soc. **76** (1970) 1310-1313.

[MP] W. Massey and F. Peterson, *The mod 2 cohomology structure of certain fibre spaces.*, Mem. Amer. Math. Soc. **74** (1967).

[MR] H. Marcum and D. Randall, *A note on self-mappings of quaternionic projective spaces*, An. Acad. Brasil. Ci. 48 (1976) 7-9.

[MG1] C. McGibbon, *Self maps of projective spaces*, Trans. Amer. Math. Soc. 271 (1982) 325-346.

[MG2] C. McGibbon, *Multiplicative properties of power maps. II*, Trans. Amer. Math. Soc. 274 (1982) 479-508.

[Mi] H. Miller, *A localization theorem in homological algebra*, Math. Proc. Camb. Phil. Soc. 84 (1978) 73-84.

[MiR] H. Miller and D. Ravenel, *Mark Mahowald's work on homotopy groups of spheres*, Contemp. Math. 146, AMS (1993) 1-30.

[Ms] G. Mislin, *The homotopy classification of self-maps of infinite quaternionic projective space*, Quart. J. Math. Oxford (2) 38 (1987) 245-257.

[Re] D. Rector, *Loop structures on the homotopy type of* S^3, Lecture Notes in Math. 249 (1971) 99-105.

[Sch] H. Scheerer, *On rationalized* H*- and co-*H*-spaces.*, Manusc. Math. **51** (1984) 63-87.

[Se] P. Selick, *Odd primary torsion in* $\pi_k S^3$, Topology 17 (1978) 407-412.

[St] J. Stasheff, H*-spaces from a homotopy point of view*, Lect. Notes in Math. **161**, Springer, Berlin (1970).

[Su] D. Sullivan, *Geometric Topology I. Localization, periodicity and Galois symmetry*, MIT Notes (1970).

[Ta] M. Tangora, *Computing the homology of the lambda algebra*, Memoirs of the Amer. Math. Soc. 337 (1985).

[Th] R. Thompson, *The* v_1*-periodic homotopy groups of an unstable sphere at odd primes*, Trans. Amer. Math. Soc. 319 (1990) 535-559.

[To] H. Toda, *Composition methods in homotopy groups of spheres*, Annals of Mathematics Studies 49, Princeton University Press (1962).

Departamento de Matemática, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisboa, Portugal
E-mail address: `ggranja@math.ist.utl.pt`